

**Your ERCP procedure is scheduled for** \_\_\_\_\_

**Please check in at:** \_\_\_\_\_ **Procedure time:** \_\_\_\_\_

**Location:**  HPI-Community Hospital North  
9800 Broadway Extension  
Oklahoma City, OK 73114

HPI-Community Hospital South  
3100 SW 89th St.  
Oklahoma City, OK 73159

## **ERCP**

Your physician has determined that ERCP is necessary for further evaluation or treatment of your condition. This brochure has been prepared to help you understand the procedure. It includes answers to questions patients ask most frequently. Please read it carefully. If you have additional questions, please feel free to discuss with the endoscopy nurse or your physician before the examination begins.

### ***What preparation is required?***

It is necessary to have a completely empty stomach for the best possible examination. You should therefore fast for at least 6 hours (and preferably overnight) before the procedure. An allergy to iodine-containing drugs (contrast material or “dye”) is not a contraindication to ERCP, but it should be discussed with your physician prior to the procedure and he should be informed of any medications you take regularly, any heart or lung conditions (or any other major diseases), and whether you have any drug allergies.

Someone must accompany you home from the procedure because of the sedation used during the examination. Even if you feel alert after the procedure, your judgment and reflexes may be impaired by your sedation for the rest of the day, making it unsafe for you to drive or operate any machinery. If a complication occurs, you may need to be hospitalized until it resolves.

### ***What is ERCP?***

ERCP is a specialized technique used to study the ducts (drainage route) of the gallbladder, pancreas, and liver (the drainage channels from the liver are called bile ducts or biliary ducts). An endoscope (flexible thin tube that allows the physician to see inside the bowel) is passed through the mouth, esophagus, and stomach into the duodenum (first part of the small intestine). After the common opening to ducts from the liver and pancreas is visually identified, a catheter (narrow plastic tube) is passed through the endoscope into the ducts. Contrast material (“dye”) is then injected gently into the ducts (pancreatic or biliary) and x-ray films are taken.

### ***Why is ERCP done?***

ERCP is a valuable tool that is used for diagnosing many diseases of the pancreas, bile ducts, liver and gallbladder. Structural abnormalities suggested by symptoms, physical examination, laboratory test or x-ray films can be shown in detail, and biopsies of abnormal tissue can be obtained if necessary. ERCP can make the important distinction between whether jaundice (yellow discoloration of eyes and skin) is caused by diseases such as hepatitis, which are treated medically, or by structural diseases, such as gallstones, tumors, or strictures (obstructing scar tissue), which are treated surgically or endoscopically. For patients who are not jaundiced but have pain or laboratory abnormalities suggesting biliary or pancreatic disease, ERCP may also provide important diagnostic information. ERCP can be used to determine whether or not surgery is necessary, and is helpful in providing the anatomic detail the surgeon needs to

# **ERCP**

plan an operation when surgery is necessary. Because x-ray films or scans may be taken, it is important that women of childbearing age tell their physician if they are pregnant. The information provided by an ERCP is far more detailed than that provided by standard x-ray films or scans.

Diagnostic ERCP is the necessary first step for therapeutic ERCP. Several conditions of the biliary or pancreatic ducts can be treated (cured or improved) by therapeutic ERCP techniques that can open the end of the bile duct, extract stones and place stents (plastic drainage tubes) across obstructed ducts to improve their drainage.

## ***What can be expected during ERCP?***

Your physician will discuss why ERCP is being performed, potential complications from ERCP and alternative diagnostic or therapeutic tests that are available. A local anesthetic may be applied to your throat and an intravenous sedative may be given to make you more comfortable during the test. Some patients also receive antibiotics before the procedure. The test begins with you lying on your left side on an x-ray table. The endoscope is passed through the mouth, esophagus, and stomach into the duodenum. The instrument does not interfere with breathing. Air is introduced through the instrument and may cause temporary bloating during and after the procedure. The injection of contrast material into the ducts rarely causes discomfort.

## ***What are possible complications of ERCP?***

ERCP is generally a well-tolerated procedure when performed by physicians who have had special training and experience in this technique. Localized irritation of the vein into which medications were given may rarely cause a tender lump that may last several weeks. The application of heat packs or hot moist towels to the area may ease discomfort.

Major complications requiring hospitalization can occur but are uncommon during diagnostic ERCP. They include serious pancreatitis and even more rarely infections, bowel perforation, and bleeding. Another potential risk of ERCP is an adverse reaction to the sedative used. The risks of the procedure vary with the indications for the test, which is found during the procedure, what therapeutic intervention is undertaken, and the presence of other major medical problems, e.g., heart or lung diseases. Your physician will tell you what is your likelihood of complications before undergoing the test.

If the therapeutic ERCP is performed (cutting an opening in the bile duct, stone removal, dilation of a stricture, stent or drain replacement, etc.), the possibility of complications is higher than with diagnostic ERCP; complications include pancreatitis, bleeding, and bowel perforation. These risks must be balanced against the potential benefits of the procedure and the risks of alternative surgical treatment of the condition. Often these complications can be managed without surgery, but occasionally they require corrective surgery.

## ***What can be expected following ERCP?***

If you are having ERCP as an outpatient, you will be kept under observation until most of the effects of the medications have worn off. Evidence of any complications of the procedure will be looked for and hospitalization may be advised if further observation is necessary. You may experience bloating or pass gas because of the air introduced during the examination. Following the procedure, you will need to stay on clear liquids and advance your diet to light bland foods. You may resume your usual diet the following day unless you are instructed otherwise.

## ***To the patient***

Because education is an important part of comprehensive medical care, you have been provided with this information to prepare you for your procedure. If you have any questions about your need for ERCP, alternative approaches to your problem, the cost of the procedure, methods of billing, or insurance coverage, do not hesitate to speak to your doctor or doctor's office staff about it. Most endoscopists are highly trained specialists and welcome your questions regarding their credentials and training. If you have questions that have not been answered, please discuss them with the endoscopy nurse or your physician before the examination begins.

